Restoration Thinning in the Cedar River Municipal Watershed

duction

ration Thinning is used in patches of small ter trees to accelerate the development of oldcharacteristics. Most Restoration Thinning in edar River Municipal Watershed occurs in the , mid-elevation Pacific silver fir (Abies amabilis) where our youngest forests are located.

gical Objectives

tain or increase tree growth rates

lerate structural development

itate future recruitment of large wood

understory light regimes

ance understory development and species

ect special habitats

ect water quality

d catastrophic losses due to large-scale ances such as fire

rescriptions

e spacing guidelines with upper diameter limit

(15' (194 tpa*) with 5" or 6" dbh** limit

pacing may vary from 11'-19'

(16' (170 tpa) with 5" or 6" dbh limit

pacing may vary from 12'-20'

guidelines and diameter limit interact with presting natural variability to yield variable density

mon tree species are retained in order to preserve ance diversity

perimental Prescriptions

felled trees into small pieces in Unit 9b to accelerate nposition and reduce fire hazard

per acre, ** diameter at breast height

ance monitoring: Are prescriptions implemented

1/50th acre plot per every 4 acres treated

mum 10 plots per Unit

eness monitoring: Do treatments achieve desired

al Objectives?

. 002-12 Treatment, 5 Control, remeasure 2007

003-16 Treatment, 5 Control, remeasure 2008 004- 15 Treatment, 5 Control, remeasure 2009



structure created by Restoration Thinning.




Figure 2. Stand density histogram for 2003 Restoration Thinning Unit 3.1. Densities were calculated from 1/50th acre plots distributed systematically throughout the unit. Note the skewed distribution, indicating variable tree




Figure 3. Forest structure and composition following Restoration Thinning in 2003 Unit 3.1. Restoration thinning preserved uncommon species, even small individuals, which would have been eliminated in a normal pre-commercial

Figure 4. Drawing of the world record Pacific silver fir by Robert Van Pelt. Restoration Thinning treatments are designed to accelerate the development of large, complex old-growth trees, such as the record Goodman Creek tree

The Restoration Thinning Progression



Figure 5. Typical pre-treatment conditions in restoration thinning units. Very high tree densities, low light environment, and virtually no understory shrubs or herbs.







Figure 10. Increment cores taken from Pacific silver fir trees thinned circa 1984. The forest from which these cores were collected lies just outside the Cedar River Watershed on US Forest Service property. The forest was thinned to approximately 12' X 12' spacing between residual trees (300 trees per acre). On average, the post-thinning radial growth rate increased by 85% (n = 14, 10 year preand post-thinning radial growth). All trees increased radial growth rate following thinning (n = 14, min = 20%, max = 160%).

lanagement

interval between thinning adjacent units to allow for down wood decomposition

de down wood treatment on all units to create fuel breaks and prevent vehicle caused ignitions and fires

sive fire patrols during periods of high fire risk

disturbance

I resources are conserved